



SAFETY APPENDIX

Table of Contents

Concussion Information	Page 1
LL Child Protection Program.....	Page 3
Pre-Game Check-List	Page 5
On-Deck Safety	Page 6
Lightening Safety	Page 7
Asthma Emergency Signs.....	Page 9
Treatment of Dental Injuries	Page 10
Warm-Up Drill Suggestions.....	Page 11
Hydration Information	Page 12



Parent/Athlete Concussion Information Sheet

A concussion is a type of traumatic brain injury that changes the way the brain normally works. A concussion is caused by bump, blow, or jolt to the head or body that causes the head and brain to move rapidly back and forth. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious.

WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury.

If an athlete reports **one or more** symptoms of concussion listed below after a bump, blow, or jolt to

Did You Know?

- Most concussions occur *without* loss of consciousness.
- Athletes who have, at any point in their lives, had a concussion have an increased risk for another concussion.
- Young children and teens are more likely to get a concussion and take longer to recover than adults.

the head or body, s/he should be kept out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it’s OK to return to play.

SIGNS OBSERVED BY COACHING STAFF	SYMPTOMS REPORTED BY ATHLETES
Appears dazed or stunned	Headache or “pressure” in head
Is confused about assignment or position	Nausea or vomiting
Forgets an instruction	Balance problems or dizziness
Is unsure of game, score, or opponent	Double or blurry vision
Moves clumsily	Sensitivity to light
Answers questions slowly	Sensitivity to noise
Loses consciousness (<i>even briefly</i>)	Feeling sluggish, hazy, foggy, or groggy
Shows mood, behavior, or personality changes	Concentration or memory problems
Can’t recall events <i>prior</i> to hit or fall	Confusion
Can’t recall events <i>after</i> hit or fall	Just not “feeling right” or “feeling down”

CONCUSSION DANGER SIGNS

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. An athlete should receive immediate medical attention if after a bump, blow, or jolt to the head or body s/he exhibits any of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that not only does not diminish, but gets worse
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people or places
- Becomes increasingly confused, restless, or agitated
- Has unusual behavior
- Loses consciousness (*even a brief loss of consciousness should be taken seriously*)

WHY SHOULD AN ATHLETE REPORT THEIR SYMPTOMS?

If an athlete has a concussion, his/her brain needs time to heal. While an athlete's brain is still healing, s/he is much more likely to have another concussion. Repeat concussions can increase the time it takes to recover. In rare cases, repeat concussions in young athletes can result in brain swelling or permanent damage to their brain. *They can even be fatal.*

It's better to miss one game than the whole season. For more information on concussions, visit: www.cdc.gov/Concussion.

Remember

Concussions affect people differently. While most athletes with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

WHAT SHOULD YOU DO IF YOU THINK YOUR ATHLETE HAS A CONCUSSION?

If you suspect that an athlete has a concussion, remove the athlete from play and seek medical attention. Do not try to judge the severity of the injury yourself. Keep the athlete out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it's OK to return to play.

Rest is key to helping an athlete recover from a concussion. Exercising or activities that involve a lot of concentration, such as studying, working on the computer, or playing video games, may cause concussion symptoms to reappear or get worse. After a concussion, returning to sports and school is a gradual process that should be carefully managed and monitored by a health care professional.

Student-Athlete Name Printed

Student-Athlete Signature

Date

Parent or Legal Guardian Printed

Parent or Legal Guardian Signature

Date



A Parent's Guide to the Little League Child Protection Program

Introduction

The backbone of Little League® is the adult volunteer. One million strong, it is this corps of dedicated people who coach the teams, umpire the games, work in the concession stands, serve on the local board of directors, and serve at the District level. These people, who live in every U.S. state and more than 100 other countries, make Little League the world's largest and most respected youth sports organization.

We know that the greatest treasure we have is children. As adults, we must ensure that these young people are able to grow up happy, healthy and, above all, safe. Whether they are our children, or the children of others, each of us has a responsibility to protect them.

The Little League Child Protection Program seeks to educate children and volunteers in ways to prevent child abusers from becoming involved in the local league. Part of that education has been to assist local Little League volunteers in finding effective and inexpensive ways to conduct background checks. Little League regulations now say: "No local league shall permit any person to participate in any manner, whose background check reveals a conviction for any crime involving or against a minor." (Reg. I [c] 9.)

Background checks were optional until the 2003 season. Effective in 2007, the local league must conduct a nationwide search that contains the applicable government sex offender registry data. Advances in computer technology – allowing greater access to public records – make it possible for background checks (at a minimum, to see if an individual is a registered sex offender in any given state) to be conducted in every U.S. state. Local Little League programs are now **required** to annually conduct a background check of Managers, Coaches, Board of Directors members and any other persons, volunteers or hired workers, who provide regular service to the league and/or have repetitive access to, or contact with, players or teams. (Reg. I [b], Reg. I [c] 9.)

The purpose of these background checks is, first and foremost, to protect children. Second, they maintain Little League as a hostile environment for those who would seek to harm children. Third, they will help to protect individuals and leagues from possible loss of personal or league assets because of litigation.

The United States Department of Justice National Sex Offender Public Registry is free and available at www.nsopr.gov.

What Can Parents Do?

Most children have been warned about the dangers of talking to strangers. But for many children, sexual molestation is committed by someone they know. In fact, 80 to 85 percent of all sexual abuse cases in the U.S. are committed by an individual familiar to the victim, according to statistics compiled by Big Brothers & Big Sisters of America.

The truth is, child sex offenders can come from every background, every occupation, every race, and every level of education. They may be married, and they may have children of their own. It is dangerous to believe that the only threat is the stranger in a long raincoat, lurking behind a tree.

In fact, the promotion of this myth may contribute to the problem. Sometimes, a child who is molested by a known and "trusted" person will feel so guilty about not reacting the "right" way that he or she never reports the problem.

Sadly, we have all seen too many reports in which teachers, police officers, clergy, youth sports volunteers, etc., trusted by all, have violated that trust and molested children in their care. Of course, this must never be tolerated in Little League or anywhere else.

In many of these situations, the young victims are actually seduced, sometimes over a period of months or even years. The child's family is lulled into believing the unusual attention being lavished is a bond of friendship between the adult and the child. In fact, the adult abuser often uses gifts, trips, attention and affection as part of a courtship process. Sometimes, the courtship process extends to the child's parent(s), but the real target is the child.

Often, but not always, the victim of this type of child sex offender is the child of a single parent. In these cases, the single parent sees the child's adult friend as a surrogate parent – a Godsend. The very opposite is true.

Two good rules of thumb for all local Little Leagues and parents

- Generally, a person involved in a local Little League program should not put himself or herself in a one-on-one situation involving a child who is not their own. Of course, some isolated situations may arise where one-on-one situations could take place. However, a one-on-one situation should not be actively *sought out* by the adult, and should not be an ongoing occurrence.

- Generally, a person involved in a local Little League program should not provide unwarranted gifts, trips, attention and affection to individual children who are not their own. The key word is *unwarranted*.

Warning Signs of a Seducer

While it remains important to teach young children about the dangers of accepting items from strangers, or talking to them, we should all beware of the danger posed by the “seducer-type” child sex offender.

Each of the individual signs below means very little. Taken as a group, however, the signs *MAY* point to this type of child sex offender, and should be applied to anyone who has repetitive access to, or contact with, children.

- Provides unwarranted gifts, trips, affection and attention to a specific child or small group of children
- Seeks access to children
- Gets along with children better than adults
- “Hangs around” children more than adults
- Has items at home or in vehicle specifically appealing to children of the ages they intend to molest, such as posters, music, videos, toys, and even alcohol or drugs
- Displays excessive interest in children (may include inviting children on camping trips or sleepovers)
- Single, over 25 years old (but could be married, sometimes as a “cover,” and could be any age)
- Photographs or videotapes children specifically
- Lives alone, or with parents
- Refers to children as objects (“angel,” “pure,” “innocent,” etc.)
- Manipulates children easily

Again, each of these items, by themselves, is relatively meaningless. Taken together, however, they may indicate a problem.

What to Watch For in Your Child

We’ve seen the signs that could point to a child sex offender, but what about the signs a child might display when he or she has been sexually abused or exploited? Some of these symptoms may be present in a child who has been or is being sexually abused, when such symptoms are not otherwise explainable: sudden mood swings, excessive crying, withdrawal, nightmares, bed-wetting, rebellious behavior, fear of particular people or places, infantile behavior, aggressive behavior, and physical signs such as pain, itch, bleeding, fluid or rawness in private areas.

Getting More Information

These items are meant solely as a general guide, and should not be used as the only means for rooting out child sex offenders. Parents can access more information on child abuse through the National Center for Missing and Exploited Children (a non-profit organization founded by John Walsh, <http://www.missingkids.com/>) and the National Clearinghouse on Child Abuse and Neglect

Information (part of a service of the Children’s Bureau, within the Administration on Children, Youth and Families, Administration for Children and Families, U.S. Department of Health and Human Services, <http://www.calib.com/nccanch/>).

How to Report

Suspected Child Maltreatment

The National Clearinghouse on Child Abuse and Neglect Information advises this: If you suspect a child is being maltreated, or if you are a child who is being maltreated, call the Childhelp USA National Child Abuse Hotline at 1-800-4-A-CHILD (1-800-422-4453; TDD [text telephone] 1-800-2-A-CHILD). This hotline is available 24 hours a day, seven days a week. The Hotline can tell you where to file your report and can help you make the report.

Or, for a list of states’ toll-free telephone numbers for reporting suspected child abuse, visit the “Resource Listings” section at this site: <http://www.calib.com/nccanch/pubs/prevenres/organizations/tollfree.cfm>, or call the Clearinghouse at 1-800-FYI-3366.

Talk to Your Kids; Listen to Your Kids

It is important that you as a parent talk frankly to your children. If a child reports sexual abuse, statistics show he or she is probably telling the truth.

Unfortunately, the sexually molested child often sees himself or herself as the one “at fault” for allowing abuse to happen. Your children **MUST** know that they can come to you with this information, and that you will support them, love them, and *believe* them.

If there is an allegation of sexual abuse of a minor, the crime should be reported immediately. These criminals who steal childhood **MUST BE STOPPED**.

This brochure was produced by Little League Baseball, Incorporated; P.O. Box 3485; Williamsport, PA 17701

Little League Baseball and Softball does not limit participation in its activities on the basis of disability, race, creed, color, national origin, gender, sexual preference or religious preference.



HAVE YOU:

- Walked field for debris/foreign objects**
- Inspected helmets, bats, catchers' gear**
- Made sure a First Aid kit is available**
- Checked conditions of fences, backstops, bases and warning track**
- Made sure a working telephone is available**
- Held a warm-up drill**

Don't Swing It

...Until You're Up to the Plate!



(Photos from North Scott, Iowa, Little League)

Don't let this happen to you, or to a teammate.

REMEMBER:

Don't pick up your bat until you leave the dugout, to approach the plate.

RULE 1.08, Notes

"1. The on-deck position is not permitted in Tee Ball, Minor League or Little League (Majors) Division. 2. Only the first batter of each half-inning will be allowed outside the dugout between the half-innings in Tee Ball, Minor League or Little League (Majors) Division."

What to do if someone is struck by lightning

- ▶ **Lightning victims do not carry an electrical charge, are safe to handle, and need immediate medical attention.**
- ▶ **Call for help.** Have someone call 9-1-1 or your local ambulance service. Medical attention is needed as quickly as possible.
- ▶ **Give first aid.** Cardiac arrest is the immediate cause of death in lightning fatalities. However, some deaths can be prevented if the victim receives the proper first aid immediately. Check the victim to see that they are breathing and have a pulse and continue to monitor the victim until help arrives. Begin CPR if necessary.
- ▶ **If possible, move the victim to a safer place.** An active thunderstorm is still dangerous. Don't let the rescuers become victims. Lightning CAN strike the same place twice.



NOAA

STAY INFORMED

Listen to NOAA Weather Radio for the latest forecast and for any severe thunderstorm WATCHES or WARNINGS. Severe thunderstorms produce winds of 58 mph or greater, or hail 3/4 of an inch or larger in diameter.

A severe thunderstorm WATCH is issued when conditions are favorable for severe weather to develop.

A severe thunderstorm WARNING is issued when severe weather is imminent. National Weather Service personnel use information from weather radar, satellite, lightning detection, spotters, and other sources to issue these warnings.



NOAA WEATHER RADIO IS THE BEST WAY TO RECEIVE FORECASTS AND WARNINGS FROM THE NATIONAL WEATHER SERVICE.

Remember that all thunderstorms produce lightning and all lightning can be deadly to those outside.

Lightning Safety Awareness Week is the last full week of June. For additional information on lightning or lightning safety, visit NOAA's lightning safety web site:

<http://www.lightningsafety.noaa.gov>

or contact us at:

**National Weather Service
P.O. Box 1208
Gray, Maine 04039**

GYX 0301 (August 2003) - Revised

Coach's and Sports Official's Guide to Lightning Safety...



NOAA

LIGHTNING...

the underrated killer!

A SAFETY GUIDE

**U.S. DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND
ATMOSPHERIC ADMINISTRATION**



**NATIONAL WEATHER
SERVICE**

Gray, Maine

This safety guide has been prepared to help coaches and sports officials recognize the dangers of lightning and take appropriate safety precautions.

LIGHTNING KILLS

Play It Safe!

Each year in the United States, more than four hundred people are struck by lightning. On average, about 70 people are killed and many others suffer permanent neurological disabilities. Most of these tragedies can be avoided if proper precautions are taken. When thunderstorms threaten, coaches and sports officials must not let the desire to start or complete an athletic activity hinder their judgment when the safety of participants and spectators is in jeopardy.

It is important for coaches and officials to know some basic facts about lightning and its dangers

- ▶ **All thunderstorms produce lightning and are dangerous.** In an average year, lightning kills more people in the U.S. than either tornadoes or hurricanes.
- ▶ **Lightning often strikes outside the area of heavy rain and may strike as far as 10 miles from any rainfall.** Many deaths from lightning occur ahead of storms because people wait too long before seeking shelter, or after storms because people return outside too soon.
- ▶ **If you hear thunder, you are in danger.** Anytime thunder is heard, the thunderstorm is close enough to pose an immediate lightning threat to your location.
- ▶ **Lightning leaves many victims with permanent disabilities.** While only a small percentage of lightning strike victims die, many survivors must learn to live with very serious, life-long disabilities.

To avoid exposing athletes and spectators to the risk of lightning take the following precautions

- ▶ **Postpone activities if thunderstorms are imminent.** Prior to an event, check the latest forecast and, when necessary, postpone activities early to avoid being caught in a dangerous situation. Stormy weather can endanger the lives of participants, staff, and spectators.
- ▶ **Plan ahead.** Have a lightning safety plan. Know where people will go for safety, and know how much time it will take for them to get there. Have specific guidelines for suspending the event or activity so that everyone has time to reach safety before the threat becomes significant. Follow the plan without exception.
- ▶ **Keep an eye on the sky.** Pay attention to weather clues that may warn of imminent danger. Look for darkening skies, flashes of lightning, or increasing wind, which may be signs of an approaching thunderstorm.
- ▶ **Listen for thunder.** If you hear thunder, immediately suspend your event and instruct everyone to get to a safe place. Substantial buildings provide the best protection. Once inside, stay off corded phones, and stay away from any wiring or plumbing. Avoid sheds, small or open shelters, dugouts, bleachers, or grandstands. If a sturdy building is not nearby, a hard-topped metal vehicle with the windows closed will offer good protection, but avoid touching any metal.

- ▶ **Avoid open areas.** Stay away from trees, towers, and utility poles. Lightning tends to strike the taller objects.
- ▶ **Stay away from metal bleachers, backstops and fences.** Lightning can travel long distances through metal.
- ▶ **Do not resume activities until 30 minutes after the last thunder was heard.**
- ▶ **As a further safety measure, officials at outdoor events may want to have a tone-alert NOAA Weather Radio.** The radio will allow you to monitor any short-term forecasts for changing weather conditions, and the tone-alert feature can automatically alert you in case a severe thunderstorm watch or warning is issued. To find your nearest NOAA weather radio transmitter, go to <http://www.nws.noaa.gov/nwr/> and click on "Station Listing and Coverage."

If you feel your hair stand on end (indicating lightning is about to strike)

- ▶ **Crouch down on the balls of your feet, put your hands over your ears, and bend your head down.** Make yourself as small a target as possible and minimize your contact with the ground.



NOAA

- ▶ **Do not lie flat on the ground.**

Asthma Emergency Signs

Seek Emergency Care If A Child Experiences Any Of The Following:

- + Child's wheezing or coughing does not improve after taking medicine (15-20 minutes for most asthma medications)
- + Child's chest or neck is pulling in while struggling to breathe
- + Child has trouble walking or talking
- + Child stops playing and cannot start again
- + Child's fingernails and/or lips turn blue or gray
- + Skin between child's ribs sucks in when breathing

Asthma is different for every person.

The "Asthma Emergency Signs" above represent general emergency situations as per the National Asthma Education and Prevention Program 1997 Expert Panel Report.


If you are at all uncertain of what to do in case of a breathing emergency...

Call 9-1-1 and the child's parent/guardian!

Michigan Asthma Steering Committee of the Michigan Department of Community Health

(From the Grandville, Mich., Little League 2001 Safety Plan)

Emergency Treatment of Athletic Dental Injuries



Professionally-made, properly fitted Custom Mouthguards greatly reduce the risk and severity of mouth injuries. Mouthguards are recommended injury prevention equipment for all at-risk sports.

AVULSION (Entire Tooth Knocked Out)

1. Avoid additional trauma to tooth while handling. **Do Not** handle tooth by the root. **Do Not** brush or scrub tooth. **Do Not** sterilize tooth.
2. If debris is on tooth, *gently* rinse with water.
3. If possible, reimplant and stabilize by biting down gently on a towel or handkerchief. Do only if athlete is alert and conscious.
4. If unable to reimplant:
 Best - Place tooth in Hank's Balanced Saline Solution, i.e. "Save-a-tooth."
 2nd best - Place tooth in milk. Cold whole milk is best, followed by cold 2% milk.
 3rd best - Wrap tooth in saline-soaked gauze.
 4th best - Place tooth under athlete's tongue. Do this **ONLY** if athlete is conscious and alert.
 5th best - Place tooth in cup of water.
5. Time is very important. Reimplantation within 30 minutes has the highest degree of success rate.
TRANSPORT IMMEDIATELY TO DENTIST.

LUXATION (Tooth in Socket, But Wrong Position)

THREE POSITIONS

EXTRUDED TOOTH - Upper tooth hangs down and/or lower tooth raised up.

1. Reposition tooth in socket using firm finger pressure.
2. Stabilize tooth by gently biting on towel or handkerchief.
3. **TRANSPORT IMMEDIATELY TO DENTIST.**

LATERAL DISPLACEMENT - Tooth pushed back or pulled forward.

1. Try to reposition tooth using finger pressure.
2. Athlete may require local anesthetic to reposition tooth; if so, stabilize tooth by gently biting on towel or handkerchief.
3. **TRANSPORT IMMEDIATELY TO DENTIST.**

INTRUDED TOOTH - Tooth pushed into gum - looks short.

1. Do nothing - avoid any repositioning of tooth.
2. **TRANSPORT IMMEDIATELY TO DENTIST.**

FRACTURE (Broken Tooth)

1. If tooth is totally broken in half, save the broken portion and bring to the dental office as described under Avulsion, Item 4. Stabilize portion of tooth left in mouth by gently biting on towel or handkerchief to control bleeding.
2. Should extreme pain occur, limit contact with other teeth, air or tongue. Pulp nerve may be exposed, which is extremely painful to athlete.
3. Save all fragments of fractured tooth as described under Avulsion, Item 4.
4. **IMMEDIATELY TRANSPORT PATIENT AND TOOTH FRAGMENTS TO DENTIST.**

Academy for Sports Dentistry
875 North Michigan Ave.
Suite 4040
Chicago, IL 60611-1901

1800-273-1788
1800-ASD-1788

The Academy for Sports Dentistry, a professional organization dedicated to the dental needs of athletes at risk to sports injuries, recommends that every sports medicine team include a dentist knowledgeable in sports dentistry.

MOUTHGUARDS SHOULD NOT BE OPTIONAL EQUIPMENT

Suggestions for Warm-up Drills



Heel Cord Stretches

Lean against a wall. Reach one leg behind you. Keep the knee straight, heel on the ground, and toes pointed forward. Slightly bend the leg that's closer to the wall. Lean forward. You should feel the stretch along the back of your calf. Repeat with other leg.



Head and Neck Circles

Make a circle with your head, going around first in one direction five times. Then reverse and make five circles in the opposite direction.



Low Back Stretches

Lie on your back, bring one knee up, and pull the knee slowly toward your chest. Hold and repeat three times. Switch legs and repeat.



Shoulder Stretches #1

Stand or sit, holding your throwing arm at the wrist with your other hand. Put your arm over your head and pull gently, feeling your upper arm against your head. You should feel the stretch inside your shoulder.



Shoulder Stretches #2

Stand or sit, holding onto the elbow of your throwing arm with your other hand. Gently pull your throwing arm across your chest. You should feel the stretch inside your shoulder, especially at the back.



Shoulder Stretches #3

Stand or sit with your pitching arm out to the side and your elbow bent. Move your arm back until you feel the stretch in the front of your shoulder.




Thigh Stretches #1

Sit on the ground. Stretch both legs out in front of you. Reach forward, touching your toes. Eventually, you want to lean forward far enough to put your head on your knees. You should feel the stretch along the backs of your legs.

Thigh Stretches #2

Sit on the ground with one leg stretched out in front of you. Bend the other knee and put your foot behind you. Lean backwards. You should feel the stretch along the front of your thigh.





**WHEN IT'S HOT,
DRINK BEFORE
YOU'RE THIRSTY.**

© 1996 Little League Baseball® and Musco Lighting, Inc.

Drinking Guidelines For Hot Day Activities

Before: Drink 8 oz. immediately before exercise

During: Drink at least 4 oz. every 20 minutes

After: Drink 16 oz. for every pound of weight lost

Dehydration signs: Fatigue, flushed skin, light-headed

What to do: Stop exercising, get out of sun, drink

Severe signs: Muscle spasms, clumsiness, delirium